

FINAL REGISTRATION REPORT

Part B

Section 0

Product Background, Regulatory Context and
GAP information

Product code: SAP09I

Product name: VOLKETE MAX

Chemical active substance:

Spinosad, 480 g/L

Central Zone

Zonal Rapporteur Member State: Poland

CORE ASSESSMENT

(New product authorization according to article 34 of Regulation (EC) 1107/2009)

Applicant(s): Albaugh TKI d.o.o.

Submission date: January 2025

MS Finalisation date: 12/2025

Version history

When	What
January 2025	V0 – Original dRR version from applicant Albaugh TKI d.o.o. for submission to zRMS Poland in the frame of a new authorization according to Article 33/34 of Regulation (EC) No. 1107/2009.
April 2025	Dossier sent for evaluation
October 2025	zRMS evaluation of dRR
November 2025	RR update with additional risk assessment
December 2025	Final RR

Table of Contents

0	Product background, regulatory context and GAP information	4
0.1	Introduction.....	4
0.1.1	Reason for application	4
0.1.2	Details of zRMS(s) and concerned MS	4
0.1.3	Regulatory history of the active(s).....	5
0.1.3.1	Spinosad.....	5
0.1.4	Regulatory history of the product (if relevant)	6
0.2	zRMS conclusion	6
Appendix 1	ALL intended uses	9

zRMS comments:

The text highlighted in grey was provided by the zRMS.

0 Product background, regulatory context and GAP information

0.1 Introduction

0.1.1 Reason for application

This application follows the data requirements for the active substance laid down in Regulation (EC) No. 283/2013 or 544/2011 and the data requirements for the plant protection product laid down in Regulation (EC) No. 284/2013 or 545/2011.

The application is for an authorisation of a new generic plant protection product VOLKETE MAX according to Regulation (EC) No. 1107/2009 in Poland. SAP09I is a SC formulation containing 480 g/L Spinosad to be used as insecticide. This application is based on the data package and assessments of the reference product Dossier of SPINTOR 480 SC; registered in Poland since 2003 (No. R-18/2003 of 12 September 2003), and currently with registration No. R-47/2018wu of 19 October 2018, pursuant article 34 of Regulation (EC) 1107/2009 after the period for data protection for SPINTOR 480 SC data package has expired.

The applicant Albaugh TKI d.o.o. is part of the task force “Spinosad 48% SC JDDA”

Members of the Task Force “Spinosad 48% SC JDDA” details:

ASCENZA PRODUCTOS PARA AGRICULTURA, S.A.U.

Parque Empresarial Táctica
c/Botiguers 3, planta 4
46980 Paterna (Valencia)
Spain

INDUSTRIAS AFRASA, S.A. (now Albaugh TKI d.o.o.)

Grajski trg 21
SI-2327 RAČE
SLOVENIA

LAINCO, S.A.

Avda. Bizet, 8-12
08191, Rubí (Barcelona)
Spain

0.1.2 Details of zRMS(s) and concerned MS

Table 0.1-1: Overview of zRMS and cMS

	zRMS, product name and authorization no. (if relevant)	(if relevant) Concerned MS, MS' product name and authorization number (if applicable)
Central zone	Poland VOLKETE MAX_not available (first registration)	-

	zRMS, product name and authorization no. (if relevant)	(if relevant) Concerned MS, MS' product name and authorization number (if applicable)
	Ireland VOLKETE MAX_registration on going	-
	Belgium VOLKETE MAX_registration on going	-
Southern zone	Greece VOLKETE MAX_14889	Portugal VOLKETE MAX_registration on going
	Portugal VOLKETE_AV02189	Greece VOLKETE_14891 Cyprus VOLKETE_4108 Bulgaria VOLKETE_registration on going Spain VOLKETE_ES-01817 Italy VOLKETE_registration on going
	Spain VOLKETE MAX_registration on going	-

0.1.3 Regulatory history of the active(s)

0.1.3.1 Spinosad

Table 0.1-2: Summary of regulatory history of CAS No: 168316-95-8

Status	
Approved in EU	Y
Original Inclusion Directive	Commission confirmed in its Decision 2000/210/EC
RMS	NL
Date of Approval (or most recent renewal) of Active Substance (date of Regulation to be applied)	01/02/2007
Current expiration of approval	15/03/2025 (to be extended) 31/10/2026
Low risk substance or Candidate for Substitution?	N

Issues that need to be considered as part of the EU approval are listed below.

In this overall assessment Member States must pay particular attention to:

- On the basis of the proposed and supported uses (as listed in Appendix II), no particular issues have been identified as requiring short term attention from the Member States

The SANCO report for Spinosad (SANCO/1428/01 – rev. final – 14/07/2006) is considered to provide the relevant information on the evaluation or a reference to where such information can be found.

Table 0.1-3: Information on minimum purity of spinosad

EU agreed minimum purity from Inclusion Directive or Implementing regulation	(if different) Minimum purity of active substance used in the product / information on available equivalency report *, **
minimum purity of active substance: 850 g/kg with 50-95% spinosyn A and 5-50% spinosyn D g/kg technical product.	minimum purity of active substance: 960 g/kg Equivalence report available: Y RMS: NL

* Since EU approval new studies on the active substance have been performed (e.g. new manufacturing site, new specification) and as a result the purity of the active substance has changed (see Part C).

** If the specification of the active substance is different to that used as reference specification for EU approval then please refer to the equivalency document from the RMS.

0.1.4 Regulatory history of the product (if relevant)

Not relevant as the product has not yet been authorised.

0.2 zRMS conclusion

Section 1, 2 and 4. Identity, physical and chemical properties and further information

The three-year shelf life is accepted.

Section 3. Efficacy

The applicant changed the GAP table after the commenting period, because of the ecotox. risk assessment and asked for re-evaluation according the changed GAP table. A new evaluation was made according the changed GAP table. Uses no.: ~~1, 2, 3, 4, 5~~ 1, 1b, 2, 2b, 5 are acceptable; Uses, 3, 3a, 4, 4a are accepted with reservation/ further restriction; uses 1a, 2a, 5a, 5b, 5c are not accepted in the GAP table of the Efficacy Section (B3).

Section 5. Analytical methods

Methods for residues were provided in the dossier for the authorization of the product SPINTOR 480 SC and they are valid for the product SAP09I.

Section 6. Mammalian Toxicology

Not classified for human health, only EUH208 - *Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.*

Operator protection (only vehicle mounted): Low vegetables – workwear, Orchards – workwear and protected hands

Worker exposure: workwear, Harvest and Maintenance (Brassica) and Orchards (Searching/reaching/picking) – gloves, Orchards (Maintenance/thinning) – gloves and re-entry period of 11 days

Resident/Bystander: Low vegetables 2-3 m buffer zone, Orchards 5-m buffer zone

Section 7. Metabolism and Residues

The application is for authorization of the plant protection product SAP09I, based on the authorised product SPINTOR 480 SC in Poland since 2003 (Registration no. No. R-18/2003 of 12 September 2003), and currently with registration No. R-47/2018wu of 19 October 2018, pursuant article 33/34 of Regulation (EC) 1107/2009 after the period for data protection for SPINTOR 480 SC data package has expired.

Head Cabbage

The data submitted shows that no exceedance of the MRL of 2 mg/kg for cabbage (STMR 0.015 mg/kg, HR 0.17 mg/kg) will occur.

The intervals between applications should correspond to the one in the residue studies, i.e. 10 days.

Cauliflower, broccoli

The data submitted shows that no exceedance of the MRL of 2 mg/kg for cauliflower (STMR 0.01 mg/kg, HR 0.02 mg/kg) will occur.

The intervals between applications should correspond to the one in the residue studies, i.e. 10 days.

According to the guideline SANTE/2019/12752 Rev.02, the extrapolation from cauliflower to broccoli is not possible. It is possible to extrapolate from 4 trials on cauliflower + 4 trials on broccoli to whole subgroup of flowering brassica. However, the applicant did not provide any residue studies for broccoli.

There is insufficient data to confirm that the applicable MRL for broccoli is not exceeded.

Onion, garlic, shallots

The data submitted shows that no exceedance of the MRL of 0.07 mg/kg for onion (STMR 0.002 mg/kg, HR 0.05 mg/kg) will occur.

The calculated using the OECD calculator indicates a higher value than the current MRL. However, the MRL in force was calculated for MRL setting, based on the same data as that provided by the applicant (EFSA Journal 2012;10(3):2630). Additionally, it should be noted that residue studies were performed with 4 applications at a dose of approximately 0.096 kg as/ha and the intended GAP applies to a max. 3 applications at this dose (max 1 appl. at dose 76.8 g as./ha in the new proposed by the applicant GAP).

According to the guideline SANTE/2019/12752 Rev.02, the extrapolation from onion to garlic and shallots is possible. The data submitted show that no exceedance of the MRL of 0.07 mg/kg for garlic and shallots will occur.

Leek

The data submitted shows that no exceedance of the MRL of 0.2 mg/kg for leek (STMR 0.065 mg/kg, HR 0.17 mg/kg) will occur.

The calculated using the OECD calculator indicates a higher value than the current MRL. However, the MRL in force was calculated for MRL setting, based on the same data as that provided by the applicant (EFSA Journal 2012;10(3):2630). Additionally, it should be noted that residue studies were performed with 4 applications at a dose of approximately 0.096 kg as/ha and the intended GAP applies to a max. 3 applications at this dose (max 1 appl. at dose 76.8 g as./ha in the new proposed by the applicant GAP).

Apple, pear, wild apple, quince

The data submitted shows that no exceedance of the MRL of 0.3 mg/kg for apple (STMR 0.04 mg/kg, HR 0.12 mg/kg) will occur.

According to the guideline SANTE/2019/12752 Rev.02, the extrapolation from apple to pear, wild apple, quince is possible. The data submitted show that no exceedance of the MRL of 0.3 mg/kg for pear, wild apple and quince will occur.

Honey

Studies indicating residue levels in crops with melliferous capacity are currently required. For the intended GAP for SAP09I this applies to apples, pears, crab apples and quinces. However, the proposed uses exclude the flowering phase (BBCH limited to: before BBCH 59 and after BBCH 70). It can therefore be concluded that for the uses proposed in the intended GAP, residue studies in honey are not required.

Estimation of the chronic and acute exposure provided in EFSA Journal 2021;19(2):6404 covers residue results adequate for this dossier.

During the first peer-review and approval of the active substance an ARfD was not deemed necessary. However, in the framework of the renewal of the approval of spinosad an ARfD of 0.1 mg/kg bw was proposed (EFSA Journal 2018;16(5):5252). It is noted that this value has not yet been formally adopted by the European Commission. Nevertheless, the proposed ARfD of 0.1 mg/kg bw was used in the estimation of the exposure. Chronic and acute exposure calculations were performed using revision 3.1 of the EFSA PRIMo. Based on these calculations, EFSA concludes that no risk to consumers was identified. Nonetheless, EFSA underlines that these conclusions were reached while considering residues of only spinosyn A and D, and it may need to be reconsidered pending revision of the residue definition for risk assessment as proposed by the peer review in the framework of the renewal.

The new GAPs proposed by the applicant are less critical than those previously assessed. No additional assessment is required.

Section 8. Environmental Fate

In accordance with proposed pattern use, an exposure assessment for the formulation of CHR/F/CYNIL 750 WG was submitted.

The mitigation measures were proposed, and final decision will be made in ecotoxicological section.

Section 9. Ecotoxicology

~~Based on the risk assessment in section of ecotoxicology it can be concluded that the proposed uses of SAP09I poses acceptable risk to non target organisms, except soil mesofauna. For *F. candida*, the risk assessment, based on laboratory study, indicated a potential for adverse effects of SAP09I proposed use pattern. Therefore, the refined PEC_{SOIL} or additional higher tier data are necessary to demonstrate the acceptable risk for soil organisms. Pending the submission of mentioned information, the risk is considered unacceptable for all proposed uses of SAP09I.~~

During the commenting process, the Applicant introduced modifications to the proposed use pattern of SAP09I (Volkete Max). Additional risk assessment was performed for mammals and soil organisms. Acceptable risk for non-target organisms was demonstrated for following uses:

- Use no. 1, 2 (cabbage, broccoli, cauliflower) application rate:
 - 3 x 0.1 L/ha (3 x 48 g a.s./ha), interval: 10d, BBCH 10-49
 - 2 x 0.16 L/ha (2 x 76.8 g a.s./ha), interval: 10d, BBCH 20-49
- Uses no. 3, 4, 7 (onion, leek, garlic) application rate:
 - 1 x 0.15-0.16 L/ha (1 x 72-76.8g a.s./ha), BBCH 40-49
- Uses no. 5, 6 (pear, wild apple, quince) application rate:
 - 1 x 0.3 L/ha (1x 144 g a.s./ha) BBCH >70
 - 2 x 0.225 L/ha (2 x 108 g a.s./ha) BBCH >70, interval: 10d
 - 2 x 0.225 L/ha (2 x 108 g a.s./ha) BBCH 40-59 (1st appl.) / BBCH>70 (2nd appl.), interval: 10d

Uses to be considered safe on the basis of EU methodology:

1b, 2b - cauliflower, 3a, 4a, 6a, 6b, 6c, 7a

Uses to be considered non-safe on the basis of EU methodology:

~~All evaluated uses~~ 1, 1a, 2, 2a, 2b – broccoli, 3, 4, 5, 5', 5a, 5b, 5c, 6, 6' 7

Appendix 1 ALL intended uses

PPP (product name/code): SAP09I / Spinosad 480 SC

Active substance 1: Spinosad

Applicant: Albaugh TKI d.o.o.

Zone(s): Central ^(d)

Verified by MS: yes

Formulation type: SC ^(a, b)

Conc. of as 1: 480 g/L ^(c)

Professional use: ☒

Non professional use: ☐

GAP rev. 1, date: 2024-Jan-01

Field of use: insecticide

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. (e)	Member state(s)	Crop and/ or situation (crop destination / purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha (f)
					Method / Kind	Timing / Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg or L product / ha a) max. rate per appl. b) max. total rate per crop/season	g as/ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min / max		
Zonal uses (field or outdoor uses)													
I	PL	Head cabbage [BRSOL]	F	<i>Pieris brassicae</i> [PIERBR] <i>Artogeia rapae</i> [PIERRA] <i>Mamestra brassicae</i> [BARABR]	Spray	Young caterpillars / first damages BBCH 10-49	a) 3-2 b) 3-2	7-10	a) 0.2 b) 0.6-0.4	a) 96 b) 288 192	2400- 600	3	Dose rate 0.1-0.2 L/ha Residues: The intervals between applications should correspond to the one in the residue studies, i.e. 10 days.

2	PL	Broccoli [BRSOK] Cauliflower [BRSOL]	F	<i>Artogeia rapae</i> [PIERRA] <i>Mamestra brassicae</i> [BARABR]	Spray	Young caterpillars / first damages BBCH 10-49	a) 3-2 b) 3-2	10	a) 0.2 b) 0.6 L/ha	a) 96 b) 288 192	2400- 600	3	Dose rate 0.1-0.2 L/ha Broccoli: Residues: N There is insufficient data to confirm that the applicable MRL for broccoli is not exceeded. Cauliflower: Residues: The intervals between applications should correspond to the one in the residue studies, i.e. 10 days.
3	PL	Onion [ALLCE]	F	<i>Thrips tabaci</i> [THRITB]	Spray	Notice of the pest / first damages BBCH 10-39	a) 3 b) 3	10	a) 0.2 L/ha b) 0.6 L/ha	a) 96 b) 288	2300- 600	7	Dose rate 0.15-0.2 L/ha
4	PL	Leek [ALLCE]	F	<i>Thrips tabaci</i> [THRITB]	Spray	Young caterpillars / first damages BBCH 10-39	a) 3 b) 3	10	a) 0.2 L/ha b) 0.6 L/ha	a) 96 b) 288	200- 600	7	Dose rate 0.15-0.2 L/ha
5	PL	Apple [MABSD]	F	<i>Adoxophyes orana</i> [CAPURE]	Spray	BBCH 569 (up to 1 treatment) >BBCH 70 (up to 2 treatments)	a) 2 b) 2	10 21	a) 0.3 L/ha b) 0.6 L/ha	a) 144 b) 288	300- 1500	14	Maximum number of treatments per growing season: 2 0.23 L/1000 m ² of LWA
Minor uses according to Article 51													
6	PL	Pear [PYUCO] Crab apple [MABSY] Quince [CYDOB]	F	<i>Adoxophyes orana</i> [CAPURE]	Spray	BBCH 40-59 (up to 1 treatment) >BBCH 70 (up to 2 treatments)	a) 2 b) 2	10 21	a) 0.3 L/ha b) 0.6 L/ha	a) 144 b) 288	300- 1500	14	Maximum number of treatments per growing season: 2 0.23 L/1000 m ² of LWA

7	PL	Garlic [ALLSA] Shallot [ALLAS]	F	<i>Thrips</i> sp. [THRISP]	Spray	Notice of the pest / first damages BBCH 10-39	a) 3 b) 3	10	a) 0.2 L/ha b) 0.6 L/ha	a) 96 b) 288	200- 600	7	Dose rate 0.15-0.2 L/ha
---	----	---	---	-------------------------------	-------	---	--------------	----	----------------------------	-----------------	-------------	---	----------------------------

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(a)	Member state(s)	Crop and/ or situation (crop destination / purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Method / Kind	Timing / Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	L product / ha a) max. rate per appl. b) max. total rate per crop/season	g as/ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min / max	PHI (days)	Remarks: e.g. g safener/synergist per ha ^(b)
Zonal uses (field or outdoor uses, certain types of protected crops)													
1	PL	Head cabbage [BRSOL]	F	<i>Pieris brassicae</i> [PIERBR] <i>Artogeia rapae</i> [PIERRA] <i>Mamestra brassicae</i> [BARABR]	Spraying	Young caterpillars / first damages BBCH 10-49	a) 2-3 b) 2-3	10	a) 0.2 b) 0.6	a) 96 b) 288	200- 600	3	Dose range 0.1-0.2 L/ha If higher tier assessment in Folsomia is not accepted please refer to Use No. 1a,b Unacceptable risk for soil organisms
1a	PL	Head cabbage [BRSOL]	F	<i>Pieris brassicae</i> [PIERBR] <i>Artogeia rapae</i> [PIERRA] <i>Mamestra brassicae</i> [BARABR]	Spraying	Young caterpillars / first damages BBCH 20-49	a) 2 b) 2	10	a) 0.16 b) 0.32	a) 76.8 b) 153.6	200- 600	3	Dose range 0.1-0.16 L/ha Intended Use just in case higher tier assessment in Folsomia is not accepted
1b	PL	Head cabbage [BRSOL]	F	<i>Pieris brassicae</i> [PIERBR] <i>Artogeia rapae</i> [PIERRA] <i>Mamestra brassicae</i> [BARABR]	Spraying	Young caterpillars / first damages BBCH 10-49	a) 2-3 b) 2-3	10	a) 0.1 b) 0.3	a) 48 b) 144	2400- 600	3	Intended Use just in case higher tier assessment in Folsomia is not accepted
2	PL	Broccoli [BRSOK] Cauliflower [BRSOB]	F	<i>Artogeia rapae</i> [PIERRA] <i>Mamestra brassicae</i> [BARABR]	Spraying	Young caterpillars / first damages BBCH 10-49	a) 2-3 b) 2-3	10	a) 0.2 b) 0.6	a) 96 b) 288	200- 600	3	Dose range 0.1-0.2 L/ha If higher tier assessment in Folsomia is not accepted please refer to Use No. 2a Unacceptable risk for soil organisms

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(a)	Member state(s)	Crop and/ or situation (crop destination / purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Method / Kind	Application Timing / Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	L product / ha a) max. rate per appl. b) max. total rate per crop/season	g as/ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min / max	PHI (days)	Remarks: e.g. g safener/synergist per ha ^(b)
2a	PL	Broccoli [BRSOK] Cauliflower [BRSOB]	F	<i>Artogeia rapae</i> [PIERRA] <i>Mamestra brassicae</i> [BARABR]	Spraying	Young caterpillars / first damages BBCH 20-49	a) 2 b) 2	10	a) 0.16 b) 0.32	a) 96-76.8 b) 288-153	200- 600	3	Dose range 0.1-0.16 L/ha Intended Use just in case higher tier assessment in Folsomia is not accepted
2b	PL	Broccoli [BRSOK] Cauliflower [BRSOB]	F	<i>Artogeia rapae</i> [PIERRA] <i>Mamestra brassicae</i> [BARABR]	Spraying	Young caterpillars / first damages BBCH 20-49	a) 2-3 b) 2-3	10	a) 0.1 b) 0.3	a) 48 b) 144	2400- 600	3	Intended Use just in case higher tier assessment in Folsomia is not accepted
3	PL	Onion [ALLCE]	F	<i>Thrips tabaci</i> [THRITB]	Spraying	Notice of the pest / first damages BBCH 10-49	a) 1 b) 1	na	a) 0.15 b) 0.15	a) 72 b) 72	200- 600	7	If higher tier assessment in Folsomia is not accepted please refer to Use No. 3a Unacceptable risk for soil organisms
3a	PL	Onion [ALLCE]	F	<i>Thrips tabaci</i> [THRITB]	Spraying	Notice of the pest / first damages BBCH 40-49	a) 1 b) 1	na	a) 0.165 b) 0.165	a) 76.8-72 b) 76.8-72	2300- 600	7	Dose range 0.15-0.16 L/ha Intended Use just in case high- er tier assessment in Folsomia is not accepted
4	PL	Leek [ALLPO]	F	<i>Thrips tabaci</i> [THRITB]	Spraying	Notice of the pest / first damages BBCH 10-49	a) 1 b) 1	na	a) 0.15 b) 0.15	a) 72 b) 72	200- 600	7	If higher tier assessment in Folsomia is not accepted please refer to Use No. 4a Unacceptable risk for soil organisms
4a	PL	Leek [ALLPO]	F	<i>Thrips tabaci</i> [THRITB]	Spraying	Notice of the pest / first damages BBCH 40-49	a) 1 b) 1	na	a) 0.165 b) 0.165	a) 76.8-72 b) 76.8-72	2300- 600	7	Dose range 0.15-0.16 L/ha Intended Use just in case high- er tier assessment in Folsomia is not accepted

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(a)	Member state(s)	Crop and/ or situation (crop destination / purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Method / Kind	Application Timing / Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	L product / ha a) max. rate per appl. b) max. total rate per crop/season	g as/ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min / max	PHI (days)	Remarks: e.g. g safener/synergist per ha ^(b)
5	PL	Apple [MABSD]	F	<i>Adoxophyes orana</i> [CAPURE]	Spraying	BBCH 40-59 (up to 1 treatment) >BBCH 70 (up to 2 treatments)	a) 2 b) 2	10-21	a) 0.3 b) 0.6	a) 144 b) 288	300- 1500	14	Dose range 0.18-0.3 L/ha If higher tier assessment in Folsomia is not accepted please refer to Use No. 5a,b,e Unacceptable risk for soil organisms
5c	PL	Apple [MABSD]	F	<i>Adoxophyes orana</i> [CAPURE]	Spraying	>BBCH 70 (up to 2 treatments)	a) 2 b) 2	10	a) 0.3 b) 0.6	a) 144 b) 288	300- 1500	14	Dose range 0.18-0.3 L/ha If higher tier assessment in Folsomia is not accepted please refer to Use No. 5a,b,e Unacceptable risk for soil organisms
5a	PL	Apple [MABSD]	F	<i>Adoxophyes orana</i> [CAPURE]	Spray	BBCH >70	a) 1 b) 1	NR	a) 0.3 b) 0.3	a) 144 b) 144	300- 1500	14	Dose range 0.18-0.3 L/ha Intended Use just in case high- er tier assessment in Folsomia is not accepted
5b	PL	Apple [MABSD]	F	<i>Adoxophyes orana</i> [CAPURE]	Spraying	>BBCH 70 (up to 2 treatments)	a) 2 b) 2	10	a) 0.225 b) 0.450	a) 108 b) 216	300- 1500	14	Dose range 0.18-0.225 Intended Use just in case high- er tier assessment in Folsomia is not accepted
5e	PL	Apple [MABSD]	F	<i>Adoxophyes orana</i> [CAPURE]	Spraying	BBCH 40-59 (up to 1 treatment) >BBCH 70 (up to 2 treatments)	a) 2 b) 2	10	a) 0.225 b) 0.450	a) 108 b) 216	300- 1500	14	Dose range 0.18-0.225 Intended Use just in case high- er tier assessment in Folsomia is not accepted

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(a)	Member state(s)	Crop and/ or situation (crop destination / purpose of crop)	F. Fn. G. Gn. Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Method / Kind	Timing / Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	L product / ha a) max. rate per appl. b) max. total rate per crop/season	g as/ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min / max	PHI (days)	Remarks: e.g. g safener/synergist per ha ^(b)
Minor uses according to Article 51 (zonal uses)													
6	PL	Pear [PYUCO] Crab apple [MABSY] Quince [CYDOB]	F	<i>Adoxophyes orana</i> [CAPURE]	Spraying	BBCH 40-59 (up to 1 treatment) >BBCH 70 (up to 2 treatments)	a) 2 b) 2	10-21	a) 0.3 b) 0.6	a) 144 b) 288	300- 1500	14	Dose range 0.18-0.3 L/ha If higher tier assessment in Folsomia is not accepted please refer to Use No. 6a,b,e Unacceptable risk for soil organisms
6 ^c	PL	Pear [PYUCO] Crab apple [MABSY] Quince [CYDOB]	F	<i>Adoxophyes orana</i> [CAPURE]	Spraying	>BBCH 70 (up to 2 treatments)	a) 2 b) 2	10	a) 0.3 b) 0.6	a) 144 b) 288	300- 1500	14	Dose range 0.18-0.3 L/ha If higher tier assessment in Folsomia is not accepted please refer to Use No. 6a,b,e Unacceptable risk for soil organisms
6a	PL	Pear [PYUCO] Crab apple [MABSY] Quince [CYDOB]	F	<i>Adoxophyes orana</i> [CAPURE]	Spray	BBCH >70	a) 1 b) 1	NR	a) 0.3 b) 0.3	a) 144 b) 144	300- 1500	14	Dose range 0.18-0.3 L/ha Intended Use just in case high- er tier assessment in Folsomia is not accepted
6b	PL	Pear [PYUCO] Crab apple [MABSY] Quince [CYDOB]	F	<i>Adoxophyes orana</i> [CAPURE]	Spray	>BBCH 70 (up to 2 treatments)	a) 2 b) 2	10	a) 0.225 b) 0.450	a) 108 b) 216	300- 1500	14	Dose range 0.18-0.225 Intended Use just in case high- er tier assessment in Folsomia is not accepted
6c	PL	Pear [PYUCO] Crab apple [MABSY] Quince [CYDOB]	F	<i>Adoxophyes orana</i> [CAPURE]	Spray	BBCH 40-59 (up to 1 treatment) >BBCH 70 (up to 2 treatments)	a) 2 b) 2	10	a) 0.225 b) 0.450	a) 108 b) 216	300- 1500	14	Dose range 0.18-0.225 Intended Use just in case high- er tier assessment in Folsomia is not accepted

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(a)	Member state(s)	Crop and/ or situation (crop destination / purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Method / Kind	Timing / Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	L product / ha a) max. rate per appl. b) max. total rate per crop/season	g as/ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min / max	PHI (days)	Remarks: e.g. g safener/synergist per ha ^(d)
7	PL	Garlic [ALLSA] Shallot [ALLAS]	F	<i>Thrips</i> sp. [THRISP]	Spraying	Notice of the pest / first damages BBCH 10-49	a) 1 b) 1	na	a) 0.15 b) 0.15	a) 72 b) 72	200- 600	7	If higher tier assessment in Folsomia is not accepted please refer to Use No. 7a Unacceptable risk for soil organisms
7a	PL	Garlic [ALLSA] Shallot [ALLAS]	F	<i>Thrips</i> sp. [THRISP]	Spraying	Notice of the pest / first damages BBCH 40-49	a) 1 b) 1	na	a) 0.16 b) 0.16	a) 76.8 b) 76.8	200- 600	7	Dose range 0.15-0.16 L/ha Intended Use just in case high- er tier assessment in Folsomia is not accepted

Remarks table heading:

(a) e.g. wettable powder (WP), emulsifiable concentrate (EC), granule (GR)
(b) Catalogue of pesticide formulation types and international coding system CropLife International Technical Monograph n°2, 6th Edition Revised May 2008
(c) g/kg or g/l

(d) Select relevant
(e) Use number(s) in accordance with the list of all intended GAPs in Part B, Section 0 should be given in column 1
(f) No authorization possible for uses where the line is highlighted in grey, Use should be crossed out when the notifier no longer supports this use.

Remarks columns:

1 Numeration necessary to allow references
2 Use official codes/nomenclatures of EU Member States
3 For crops, the EU and Codex classifications (both) should be used; when relevant, the use situation should be described (e.g. fumigation of a structure)
4 F: professional field use, Fn: non-professional field use, Fpn: professional and non-professional field use, G: professional greenhouse use, Gn: non-professional greenhouse use, Gpn: professional and non-professional greenhouse use, I: indoor application
5 Scientific names and EPPO-Codes of target pests/diseases/ weeds or, when relevant, the common names of the pest groups (e.g. biting and sucking insects, soil born insects, foliar fungi, weeds) and the developmental stages of the pests and pest groups at the moment of application must be named.
6 Method, e.g. high volume spraying, low volume spraying, spreading, dusting, drench Kind, e.g. overall, broadcast, aerial spraying, row, individual plant, between the plants - type of equipment used must be indicated.

7 Growth stage at first and last treatment (BBCH Monograph, Growth Stages of Plants, 1997, Blackwell, ISBN 3-8263-3152-4), including where relevant, information on season at time of application
8 The maximum number of application possible under practical conditions of use must be provided.
9 Minimum interval (in days) between applications of the same product
10 For specific uses other specifications might be possible, e.g.: g/m³ in case of fumigation of empty rooms. See also EPPO-Guideline PP 1/239 Dose expression for plant protection products.
11 The dimension (g, kg) must be clearly specified. (Maximum) dose of a.s. per treatment (usually g, kg or L product / ha).
12 If water volume range depends on application equipments (e.g. ULVA or LVA) it should be mentioned under “application: method/kind”.
13 PHI - minimum pre-harvest interval
14 Remarks may include: Extent of use/economic importance/restrictions